

## **MODIFYING EFFECT OF HYPOKINETIC STRESS ON CHANGE PAIN SENSITIVITY OF RAT (Part 1)**

*Chuyan E. N., Zayachnikova T. V., Ravaeva M. Yu., Mironyuk I. S., Birukova E. A.*

*V. I. Vernadsky Crimean Federal University, Simferopol, Crimea, Russian Federation  
E-mail: m-ravaeva@rambler.ru*

Modifying action of a hypokinetic stress is investigated on change of behavioral phenomena in rats with experimentally induced tonic somatic, visceral, acute heat pain and electrical stimulation.

It is shown that at adaptation of rats to short hypokinetic to a stress (the first – the sixth days) increase in resistance to painful factors is noted that is expressed in reduction of duration of painful reactions, and long restriction of mobility (the seventh – the ninth days) leads to reduction of resistance to painful factors that is shown in increase in duration of painful reactions.

Thus, the hypokinetic stress changes behavioural reactions at rats as at action of painful stresses (in "formalin" and "acetic" tests), and painful sensitivity at the threshold level in tests of "a hot plate" and electrostimulation that demonstrates the modifying action of a gipokineziya on painful sensitivity of animals regardless of the nature of a painful irritant.

However, the modifying effect of a hypokinetic stress on painful sensitivity at rats in experimental painful tests depends on mobility restriction duration. At the same time change of painful sensitivity (reduction and increase) of animals at a hypokinetic stress can serve as criterion of transition of an eustress to a distress.

**Keywords:** hypokinetic stress, behavioral reactions, tonic pain, visceral pain, acute thermal pain, electrical stimulation.

### References

1. Kalyuzhnyy L. V. *The physiological mechanisms of regulation of pain sensitivity*, 216. (M.: Meditsina (1984).
2. Bragin E. O. *The neurochemical mechanisms of regulation of pain sensitivity*, 248. (M.: Izd-vo Un-ta druzhby narodov, 1991).
3. Veyn A. M., Avrutskiy M. Ya. Pain and pain relief, 280 (Meditsina, 1997).
4. Kovalenko E. A., Gurovskiy N. N. *Hypokinesia*, 307. (M.: Meditsina, 1980).
5. Chuyan E. N. Influence of millimeter waves on the development of non-thermal intensity hypokinetic stress in rats with various individual characteristics : Avtoref. dis. kand biol.nauk., 25. (Simferopol', 1992).
6. Santana Vega L. The part of individual characteristics of physical activity in the development hypokinetic stress in rats: Avtoref. dis. kand. biol. nauk: 03.00.13 /SGU, 21. (Simferopol,1991).
7. Dubuisson D. *The formalin test: a quantitative study of the analgesic effects of morphine, meperidine and brainstem stimulation in rats and cats*, 161. (1997).
8. Koster R. Acetic acid for analgesic screening, *De Beer Fed.*, 412 (1959).
9. Certificate in computer software processing, registration and automated analysis duration and frequency of different types of behaviors in animals: № 19243 vid 18.01.2007 , M. Lutsyuk, E. Dzheldubaeva.
10. O'Callaghan J. Quantification of the analgesic activity of narcotic antagonists by a modified hot-plate procedure, *Pharmacol. Exp. Ther.*, 497 (1979).
11. Papin A. A., Petrov V., Kakurin F. F. The research component of the analgesic sedation by heat sensometrii, *Anesteziologiya i reanimatologiya*, 1, 18 (1983).
12. Vasilenko A. M. Tenzoalgotometriya, *Bol' i ee lechenie*, 6, 56 (1997).
13. Slonim A. D. *Types and forms of adaptive behavior of animals: Rukovodstvo po fiziologii "Fiziologiya povedeniya. Neyrofiziologicheskie zakonomernosti"*. 23. (M: Nauka 1986).
14. Golombek D. A., Escolar E., Burin L. J. et al. Time-dependent melatonin analgesia in mice: inhibition by opiate or benzodiazepine antagonist, *Eur. J. Pharmacol.*, 1, 25. (1991)